


MICHIGAN STATE HEALTH DEPARTMENT

SUMMARY OF LOCAL CASE DATA WITH COMPARISON TO NATIONAL CONTROLS

3,412 potential participants accessed the survey, 3,241 consented, and 3,013 met eligibility criteria between 6/28 and 7/15.

MOST COMMON REASONS FOR TESTING



INDIVIDUAL PREVENTION STRATEGIES



Face Covering

30.6% of local cases
always/often wore surgical or n95 masks

National Controls (n=3,028)
37% of controls always/often wore surgical or n95 masks



Physical Distancing

45.4% of local cases
always/often avoided crowded places/events

National Controls (n=3,028)
43% of controls always/often avoided crowded places/events

TRUSTED SOURCES OF INFORMATION

Michigan Cases


#1 CDC (77.0%)

#2 My healthcare provider (75.0%)


#3 State or County health departments (66.9%)

VACCINATIONS

Plans for Child Vaccination



Info Needed for Child Vaccination Decision



SUMMARY

Most individuals (79%) are testing as a result of symptoms. 36% tested as a result of close contact.

77% of cases had at least 3 vaccination doses and 21% had a history of prior COVID infection.

45% used both PCR and rapid testing to confirm positive COVID cases. 25% used rapid testing alone while 30% used PCR.

12% have delayed or avoided medical care in the past 3 months due to concerns related to COVID-19.

15% screened positive for anxiety and 8% screened positive for depression.