

MICHIGAN STATE DEPARTMENT OF HEALTH

SUMMARY OF LOCAL CASE DATA WITH COMPARISON TO NATIONAL CONTROLS

4,153 potential participants accessed the survey, 3,813 consented, and 3,274 met eligibility criteria between 5/13 and 5/25.

MOST COMMON REASONS FOR TESTING

Symptomatic	81%
Close Contact	41%
Preparing to Travel	8%

INDIVIDUAL PREVENTION STRATEGIES



Face Covering

34% of case

always/often wore surgical or n95 masks

National Controls (n=3,028).

37% of controls always/often wore surgical or n95 masks



Physical Distancing

51% of cases

always/often avoided crowded places/events

National Controls (n=3,028).

43% of controls always/often avoided crowded places/events

TRUSTED SOURCES OF INFORMATION

MICHIGAN CASES


#1 CDC (79%)

#2 PCP (77%)


#3 Health Depart. (69%)

VACCINATIONS

Plans for Child Vaccination



Info Needed for Child Vaccination Decision



IMPLICATIONS

Most individuals (81%) are testing as a result of symptoms. 40% tested as a result of close contact.

Individual prevention strategies were similar between cases and controls.

7 in 10 cases in Michigan are likely to vaccinate their children.

Most Trusted Sources of Info: CDC, PCP, and health department